



		Middle	School	-	Comparison of Two Rates			
Indicator	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference	
Smoked cigarettes on one or more of the past 30 days	2.9	(0.4-5.4)	2.3	(2.1-2.5)			+	
Smoked cigars on one or more of the past 30 days	1.5	(0.1-2.9)	2.9	(2.6-3.2)			+	
Used smokeless tobacco on one or more of the past 30 days	4.8	(2.1-7.5)	2.1	(1.9-2.3)			+	
Used electronic cigarettes on one or more of the past 30 days	2.4	(0.4-4.5)	4.0	(3.6-4.3)			+	
Used hookah on one or more of the past 30 days	1.1	(0.0-2.3)	3.5	(3.1-3.9)		+		
Used any form of tobacco on one or more of the past 30 days*	6.9	(3.8-10.1)	5.1	(4.7-5.4)			+	
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	8.7	(4.8-12.5)	8.1	(7.6-8.6)			+	
Using electronic cigarettes is less harmful than smoking cigarettes	65.6	(51.4-79.7)	64.7	(63.2-66.1)			+	
Using hookah is less harmful than smoking cigarettes	8.5	(2.4-14.7)	42.0	(39.9-44.0)		+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	74.0	(67.5-80.5)	72.6	(71.7-73.5)			+	
Exposed to secondhand smoke during the past 7 days (in a room or car)	45.0	(40.2-49.8)	35.9	(35.0-36.7)	+			
Smoking is allowed in the home	13.5	(9.3-17.8)	8.8	(8.3-9.3)			+	

^{*}This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.





		Middle	School		Comparison of Two Rates		
Indicator	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Students who were taught about tobacco use in any classes this year	77.1	(70.3-83.9)	47.1	(45.7-48.5)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	67.8	(60.9-74.7)	67.5	(66.7-68.3)			+
Served as an active member of SWAT	7.3	(4.3-10.3)	2.9	(2.7- 3.2)	+		
Definitely did not think that smokers have more friends	43.0	(37.2-48.9)	31.7	(30.8-32.5)	+	+	
Definitely did not think that smoking helps young people "look cool" or "fit in"	82.5	(76.5-88.5)	70.2	(69.4-70.9)	+		
Lifetime asthma	19.2	(14.1-24.3)	20.2	(19.5-20.9)			+
Current asthma	11.6	(7.6-15.7)	12.1	(11.5-12.6)			+
Asthma attack in past year	12.9	(3.8-22.0)	20.0	(18.5-21.5)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	12.1	(7.5-16.7)	12.4	(11.7-13.2)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	14.9	(9.0-20.9)	16.3	(15.6-17.0)			+
Physically active for at least 60 minutes per day during the past 7 days	33.6	(25.7-41.4)	24.7	(24.0-25.6)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	40.5	(35.8-45.2)	44.6	(43.8-45.5)			+
Described themselves as slightly or very overweight	30.2	(25.2-35.2)	30.1	(29.3-30.8)			+





	High School				Comparison of Two Rates			
Indicator	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference	
Smoked cigarettes on one or more of the past 30 days	15.7	(11.0-20.4)	7.5	(7.1-7.9)	+			
Smoked cigars on one or more of the past 30 days	9.9	(4.9-14.8)	9.1	(8.6-9.6)			+	
Used smokeless tobacco on one or more of the past 30 days	12.2	(7.6-16.8)	5.4	(5.0-5.8)	+			
Used electronic cigarettes on one or more of the past 30 days	9.8	(6.0-13.5)	10.8	(10.3-11.4)			+	
Used hookah on one or more of the past 30 days	9.4	(4.6-14.3)	11.6	(11.0-12.1)			+	
Used any form of tobacco on one or more of the past 30 days*	23.9	(17.6-30.2)	14.8	(14.2-15.5)	+			
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	25.2	(18.4-31.9)	22.6	(21.8-23.3)			+	
Using electronic cigarettes is less harmful than smoking cigarettes	68.2	(60.8-75.6)	71.1	(69.9-72.3)			+	
Using hookah is less harmful than smoking cigarettes	36.2	(26.6-45.8)	57.3	(56.0-58.6)		+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	41.3	(33.9-48.8)	60.7	(59.8-61.6)		+		
Exposed to secondhand smoke during the past 7 days (in a room or car)	53.6	(48.4-58.8)	39.7	(38.8-40.6)	+			
Smoking is allowed in the home	19.7	(14.8-24.7)	8.8	(8.3-9.3)	+			

^{*}This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.





		High S	chool		Comparison of Two Rates		
Indicator	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Students who were taught about tobacco use in any classes this year	31.6	(23.8-39.3)	29.6	(28.4-30.7)			+
Parent or guardian has talked about the dangers of tobacco in the past year	50.6	(44.7-56.5)	56.7	(55.8-57.5)			+
Served as an active member of SWAT	4.9	(1.5-8.2)	3.1	(2.8- 3.4)			+
Definitely did not think that smokers have more friends	25.5	(19.6-31.3)	25.3	(24.6-26.1)			+
Definitely did not think that smoking helps young people "look cool" or "fit in"	62.3	(55.5-69.1)	67.1	(66.3-67.9)			+
Lifetime asthma	24.5	(20.6-28.4)	21.2	(20.5-21.9)			+
Current asthma	10.9	(6.8-15.0)	10.8	(10.2-11.3)			+
Asthma attack in past year	14.5	(4.8-24.2)	16.5	(15.2-17.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	12.3	(7.4-17.3)	12.3	(11.7-12.9)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	17.5	(12.7-22.3)	15.5	(14.9-16.1)			+
Physically active for at least 60 minutes per day during the past 7 days	27.6	(22.1-33.0)	21.5	(20.8-22.2)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	42.9	(38.9-46.8)	42.5	(41.7-43.4)			+
Described themselves as slightly or very overweight	34.0	(30.0-38.0)	31.2	(30.4-31.9)			+





		Youth (Ag	es 11-1	7)	Comparison of Two Rates			
Indicator	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference	
Smoked cigarettes on one or more of the past 30 days	8.5	(5.2-11.8)	4.3	(4.1-4.6)	+			
Smoked cigars on one or more of the past 30 days	5.2	(2.7-7.7)	5.4	(5.2-5.7)			+	
Used smokeless tobacco on one or more of the past 30 days	7.8	(5.0-10.7)	3.3	(3.1-3.5)	+			
Used electronic cigarettes on one or more of the past 30 days	4.9	(2.5-7.4)	7.2	(6.8-7.5)			+	
Used hookah on one or more of the past 30 days	3.7	(1.7-5.6)	7.1	(6.7-7.5)		+		
Used any form of tobacco on one or more of the past 30 days*	14.1	(10.0-18.2)	9.2	(8.8-9.6)	+			
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	15.8	(11.4-20.3)	14.8	(14.2-15.3)			+	
Using electronic cigarettes is less harmful than smoking cigarettes	65.7	(58.0-73.4)	69.4	(68.4-70.3)			+	
Using hookah is less harmful than smoking cigarettes	22.2	(15.3-29.1)	52.8	(51.7-54.0)		+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	59.8	(53.3-66.3)	67.1	(66.4-67.7)		+		
Exposed to secondhand smoke during the past 7 days (in a room or car)	48.1	(44.1-52.2)	37.5	(36.9-38.2)	+			
Smoking is allowed in the home	15.7	(12.4-19.1)	8.4	(8.0-8.7)	+			

^{*}This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.





		Youth (Ag	es 11-1	7)	Comparison of Two Rates			
Indicator	County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference	
Students who were taught about tobacco use in any classes this year	53.4	(45.1-61.7)	38.0	(37.1-39.0)	+			
Parent or guardian has talked about the dangers of tobacco in the past year	59.6	(54.4-64.7)	62.4	(61.8-63.0)			+	
Served as an active member of SWAT	6.1	(3.9-8.3)	2.8	(2.7- 3.0)	+			
Definitely did not think that smokers have more friends	34.3	(29.7-38.9)	28.2	(27.6-28.8)	+			
Definitely did not think that smoking helps young people "look cool" or "fit in"	73.1	(67.5-78.7)	68.8	(68.2-69.3)			+	
Lifetime asthma	22.6	(19.0-26.2)	20.8	(20.3-21.3)			+	
Current asthma	11.5	(8.3-14.6)	11.5	(11.1-11.9)			+	
Asthma attack in past year	13.6	(6.6-20.5)	17.7	(16.7-18.7)			+	
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	12.4	(8.9-15.9)	12.2	(11.8-12.7)			+	
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	16.3	(12.3-20.3)	15.9	(15.4-16.4)			+	
Physically active for at least 60 minutes per day during the past 7 days	30.9	(25.8-36.1)	22.7	(22.2-23.3)	+			
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.2	(37.9-44.5)	43.7	(43.1-44.4)			+	
Described themselves as slightly or very overweight	32.5	(29.2-35.9)	30.7	(30.2-31.3)			+	